


Fig



It is called as *Ficus carica* in latin; its botanical family is Moraceae (mulberry family); it is a flowering plant famous for its fruit, its fruit has many health benefits, its fruit is called as fig; please read lesson no. 28 Fig page no. 12 in my part 2 book; it is mentioned in Holy Quran chapter no. 95 Teen verse no. 1 the chapter is named as Teen which means Fig. It is also mentioned in Hadith of Ibn Abu Bakr Al Jauzi, Kanzul-ummal, Riyad As-sulihin, Bukhari; please refer my part 2 lesson no. 28 Fig. Fig is a false fruit or multiple fruit in which the flower & seeds grow together to form a single mass; depending on the type of fig-each fig can contain thousands of seeds in it; in Hadith it is mentioned as fruit of Paradise & cure for arthritis, piles.

Please visit my website www.tib-e.nabi-for-you.com for detail Islamic study on fig.

• **NAMES:**

1. It is called Teen (التين) in Quran, Hadees & Arabic.
2. Fig in English.
3. In Latin it is called *Ficus carica* Linn.
4. In Persian, Urdu & Hindi it is called as Injeer.

It is mentioned in following books of Hadith (reference are also given as Hadith number) Bukhari : 767; Ibn Abu Bakr Al-Jauzi/At-tibb Al Nabawi Harfutta; Kanzul ummal : 28280; Riyad as-Salihin : 1006;

• **Fig tree: -**



It is grown since ancient time; it is native to Middle East & western Asia, nowadays grown throughout the world to obtain its fruit & as an ornament plant; its tree is called as *Ficus carica* (common fig) & belongs to Moraceae family, it is a flowering plant & a deciduous; it grows to a height of 7-10 meters (23-33 feet) with smooth white bark. It is gynodioecious shrub (female flowers are present on one plant & hermaphrodite flower on other plant of the same species). (Hermaphrodite is part of plant that has complete or partial reproductive organs & produces gametes). It grows in dry & sunny region, in rocky area in deep & fresh soil, can also can also grown on nutritional soil; it produces two crops each year, 1st is eaten fresh (fig) & 2nd crop is dried in winter (fig) but many types of fig tree produces fruit once a year only. Its branches are muscular & twisted, widely spread & are multiple branched; its wood is weak & decay rapidly; the trunk often has large nodal tumours; the twigs are terete (slender & smooth) & pithy rather than woody. The sap contains milk latex & is irritating to human skin.

- **Leaves: -**



Leaves are bright green, single, alternative & large (up to 1 foot), they are more less deep lobed with 1-5 sinuses, rough hairy on upper side & soft hairy on the underside.

- **Flowers: -**



The tiny flowers are out of sight, clustered inside the green fruit (technically called syconium). Pollinating insects gain access in the flower through an opening at the apex of the syconium. Its flowers are mainly female & need pollination.

- **Fruit (fig): -**



Its fruit is called as Fig, figs crop twice a year; 1st crop is called as breba crop, it occurs at spring. In cold climate the breba crop is often destroyed by spring-frost, it mainly sold fresh & eaten fresh; the 2nd crop is called as main crop, is borne in the fall on the new growth. The fruiting cycle is of 120-150days.

The matured fig has tough skin (peel) of pure green or green suffused with brown or brown purple colour; the skin often cracks upon ripen & inside pulp is exposed. The interior of fig is white inner rind containing many seeds, mass bound with jelly like flesh. The whole fruit is edible including the seeds; the fruit is delicious & many health benefits & vitamins, minerals & etc; the fruits can be eaten fresh or dried.

- **Harvesting of figs: -**

Figs must be allowed to ripe fully on tree then picked up; it will not ripe if picked up when immature. A ripen fruit will be soft & start to bend at its neck part. Harvesting should be done


gently; fresh figs do not keep well & so can be stored in refrigerator only 2 to 3 days only. It is more delicious when eaten dried; it takes 4 to 5 days in sun to get dried (also 10 to 12 hours in dehydrator process & 2 to 3 days on top of car dash board); dried figs can be stored for more than 6 to 8 months.

- **pH of Natural fig is:** - pH is 4.6 or little more; it is acidic because its pH is less than 7.

pH is a measure of hydrogen ion concentration, a measure of the acidity or alkalinity of a solution. The pH scale usually ranges from 0 to 14. Aqueous solutions at 25°C with a pH less than 7 are acidic, while those with a pH greater than 7 are basic or alkaline & 7 is neutral; only aqueous solutions have pH levels, vegetable oil has no pH value. Likewise, other oils such as animal and petrochemical oils also have no pH value. Fatty acids are organic molecules often found in foods, including vegetable oils.

The pH of pure water is 7. In general, water with a pH lower than 7 is considered acidic, and with a pH greater than 7 is considered alkaline. The normal range for pH in surface water systems is 6.5 to 8.5, and the pH range for groundwater systems is between 6 and 8.5. We can add normal water to reduce the acidity.

It is Sunnat of Prophet Muhammed (s.a.w) to mixe acidic with Alkaline to make it neutral or less acidic that why He use eat dates with watermelon or cucumber or dry dates with little butter; so you can mix one acidic with alkaline; also it is Sunnat to drink honey mixed in water; also dates or raisins soaked in water over night & drink the syrup (sharbat). Remember do not soak dates & raisin together at one time; soak at separate time & drink.



- **Calories of fig:** -

100 grams of dried fig contain 209 calories & fresh contain only 43 calories.

Glycemic index & Glycemic load of fig: -

The glycemic index of fresh figs is 51 & of dried is 61; which fall under low GI category, while the glycemic load of ¼th cup (50 grams) of chopped fresh figs is around 2 & 16 of dried fig in 60 grams, fresh fig falls in low GL & dried fig falls in moderate GL.

A food is considered to have a low Glycemic index (GI) if it is 55 or less; mid-range GI if 56 to 69 & high GI if 70 or more. Glycemic index is a number. It gives you an idea about how fast your body converts the carbs in a food into glucose.

A low Glycemic load (GL) is between 1 and 10; a moderate GL is 11 to 19; and a high GL is 20 or higher. For those with diabetes, you want your diet to have GL values as low as possible.

The glycemic load (GL) of food is a number that estimates how much the food will raise a person's blood glucose level after eating it. Glycemic load accounts for how much carbohydrate is in the food and how much each gram of carbohydrate in the food raises blood glucose levels.

- **Gross health benefit & indications of dried & fresh figs: -**

It is very helpful in piles, constipation, controls blood pressure, boost health, beneficial in recovery from malaria, dengue, typhoid, weakness; controls weight because it is low in calories, strengthen the bone, joints, muscles etc; very beneficial in osteoarthritis, skin, complex, nails, hair, it is an antioxidant, prevent cancer, free radicals, removes harmful effect of UV sun rays, improves digestion, prevents colon, breast & many types of cancers, reduces risk of heart diseases & cardiovascular disease, helps liver to function, reduces cholesterol & triglycerides; helpful in anaemia, reduces blood sugar, increases libido in both male & female, helpful in treating asthma; prevents venereal diseases, reduces throat pain, prevent eye & eye sight, good for complexion & skin health, prevents ageing etc.

Clinical pharmacology: -

The petroleum ether extract from leaves of Fig is hepato-protective; beta sitosterol present in fig is anti cancer, fig has many antioxidant compounds which protect from various types of cancer.

The methanol extract of fig are antibacterial & antifungal; Hexane is antimicrobial; these also act on total inhibition against *Candida albicans* (100%) at a concentration of 500 μ g/mL and a negative effect against *Cryptococcus neoformans*; methanolic extract (75%) strongly inhibited *Microsporum canis* and ethyl acetate extract at a concentration of 750 μ g/mL.

The 80% methanol extract from the leaves of *F. carica* has been screened against *Mycobacterium tuberculosis* means Tuberculosis infection.

For Nematicidal Activity Forty different plant species were screened for their nematicidal activity against the nematodes *Bursaphelenchus xylophilus*, *Panagrellus redivivus*, and *Caenorhabditis elegans*. The leaf extract of fig showed the strongest nematicidal activity as 74.3, 96.2, and 98.4% mortality, respectively, within 72 hrs. More you study from separately explained in contents below.

- **Modern uses of fig: -**

Please note it is Sunnah to eat two opposite quality stuff together, one of hot potency (acidic) & other cold potency (alkaline).

For complexion: -

Take one fresh or dried fig soak it overnight in one glass water or cow milk, filter & drink the water early morning empty stomach & take the pulp of fig and apply on face, keep it for 20 minutes & wash the face with warm water, do it once or twice a week; by it acne reduces & heals, tightens the facial skin, clears the pores, reduces sebum & oil, improves fairness.

For constipation: -

Take 1 dried fig, soak in 1 glass of water overnight, drink & eat the pulp early morning empty stomach daily for 40 days & eat 1 teaspoon of senna powder at night & drink 1 glass water on it for 40 days then alternative days for 40 days then twice a week for 7 weeks.

For irritable bowel syndrome (IBS) or other gastric problem: -

Take 1 dried or fresh fig, 1 small size cucumber, 1 carrot, half beetroot, 3 dates, one teaspoon of barley flour roasted, 1 cup warm water, 7 seeds of black caraway (kalonji)-prepare juice out of these & drink it in breakfast for 21 days & alternative days till complete relief after relief twice a week for 21 weeks followed by once a week for 21 days.

For general health: -

Take 1 dried fig, 3 pieces of watermelon, 3 dates, 1 small cucumber, 3 almonds, 1 cup cow milk eat these at breakfast twice a week for 11 weeks followed by off for 3 weeks & again start, repeat the cycle regularly.

For heart health: -

Take 1 dried or fresh fig, 3 dates, 1 beetroot, 1 teaspoon of extra virgin olive oil, 1 small tomato, 3 pieces of watermelon prepare juice out of these & drink at breakfast 2 times a week regularly.

For bones health & to prevent cancer: -

Take 1 dried fig soak it in 1 table spoon of extra virgin olive oil overnight, put 1 table spoon of pure honey, take 3 pieces of watermelon, 3 dates, half orange eat these at breakfast for 3 times a week for 40 weeks.

For malaria, dengue & typhoid: -

Eat fresh or dried fig 3, watermelon 3 slice each time for 2 to 3 times a day for 5 days.

For piles: -

Eat 3 dried fig or fresh fig, 7 pieces of watermelon, 3 dates daily at night for 11 days followed by 11 alternative days followed by once a week for life long.

For arthritis: -

Eat 1 dried fig, 1 date, lick 1 spoon pure honey & 1 spoon aloe vera gel daily morning for 40 days & 1 cup cow milk with 1 teaspoon of turmeric powder at night for 40 days.

• **Contents/constituents of dry & fresh figs: -**

All contents may not present in all types of it, because there are many varieties of it according to geographical regions & content may differ a lot as per cultivation, soil, seed, climate etc.

It contains water 39% fresh fig, soluble fibre, potassium, calcium (fresh 35 mg & dried 162 mg), magnesium, iron, copper, vitamin A, K, E, C, prebiotic, oxalates, natural sugar (dried has 48 grams of sugar in 100 gram, fresh fig has 16 gram of sugar in 100 grams), sodium, protein, iron, vitamin B1, B2, B5, B6, folate, choline, magnesium, phosphorus, zinc, carbohydrate, carotenoid, selenium, chloride, lutein & zeaxanthin etc.

The above ingredients are based on scientific study, means these has been indentified, known & learnt by modern science, it does not means that it contains only these ingredients, there may be many more ingredients which are yet to be discovered, learnt & known by modern science.

A good quality dried fig contains following amino acids: -

Active ingredient of Fig: - phenolic compounds, phytosterols, organic acids, anthocyanin composition, triterpenoids, coumarin, and volatile compounds such as hydrocarbons.

Phenolic acids such as 3-O- and 5-O-caffeoylquinic acids, ferulic acid, quercetin-3-O-glucoside, quercetin-3-O-rutinoside and organic acids (oxalic, citric, malic, quinic) have been isolated from the water extract of the leaves of Fig.

Amino acids	Dried fig
Weight (g)	149
Tryptophan(mg) (% RDI)	30 (11%)
Threonine(mg) (% RDI)	127 (12%)
Isoleucine(mg) (% RDI)	133 (9%)
Leucine(mg) (% RDI)	191 (7%)
Lysine(mg) (% RDI)	131 (6%)
Methionine(mg) (% RDI)	51 (7%)
Cystine(mg)	54

(% RDI)	(19%)
Phenylalanine(mg) (% RDI)	113 (13%)
Tyrosine(mg) (% RDI)	61 (7%)
Valine(mg) (% RDI)	182 (10%)
Histidine(mg) (% RDI)	55 (8%)
Arginine(mg)	115
Alanine(mg)	200
Aspartic acid(mg)	961
Betaine(mg)	1
Glutamic acid(mg)	440
Glycine(mg)	161
Proline(mg)	909
Serine(mg)	191

Natural basic pharmacology of figs based on human intake in natural form: -

• **Natural oxalate: -**

It is present in many types plant, it is an organic compound found in leafy vegetables, fruits, nuts, seeds etc. In plants it usually bounds to minerals forming oxalate. The term oxalic acid & oxalate are used interchangeable in nutritional science.

Main sources of natural oxalate: -

It is present is beetroot leaves (chard leaves), beetroot, carrot, spinach, sweet potato, turnip leaves, potato, apple, strawberry, green beans, broccoli, apricots, fig etc.

Basic pharmacokinetics of oxalate (bases on human intake in natural food products): -

In the human body some of the eaten oxalate is broken down by bacteria before it combines (mainly) with calcium & iron in colon & in kidneys (also in urinary tract) to form crystals & is excreted in urine as minute crystals (little is excreted in stool also) if taken in large quantity can cause kidney stones.

Also when fat is not absorbed properly, the fats binds with calcium and leaves oxalate behind, this oxalate is taken up by kidneys to excrete it; it also prevents absorption of calcium & other minerals; Presence of prebiotic can effect oxalic acid excretion, fig contains prebiotic & oxalates both. Our body makes oxalates as an end product of protein, vitamin c etc metabolism. It is excreted in urine & stool. Oxalate present in food is soluble & insoluble, insoluble bind with calcium or other molecules that makes it much harder to absorb; soluble do not bind with other molecules & is easily absorbed. Absorption of it differs among people, for some, oxalate is largely broken down in the intestines & eliminate without causing issues; in some, large amount of oxalate is absorbed; & in some, leaky gut can increase the absorption of it & the needle shaped oxalate crystals can perforate the mucus membrane damaging the cells of intestine & cause more leakiness.

Leaky gut is also called as increased intestinal permeability, it is a condition in which many things can pass through the intestinal wall, and this occurs due to breach in intestinal wall or damaged cells, things can pass through their gasps (breach).

Excessive oxalate accumulates in crystal form in many parts of the body like joints, muscles, kidneys etc.

Basic clinical pharmacology of oxalate: -

It reduces absorption of some minerals because it binds with some mineral in the intestine, if taken in larger amount causes kidney stones, autism, vaginal pain etc; we should drink lot of water to avoid excessive oxalate collection in the body & to flush out; its benefits in human body is under research.

• **Prebiotic: -**

It is non digestible (fibre or carbohydrate) food ingredient that promotes the growth of beneficial bacterias & fungus in the intestines. It induces the growth & activities of helpful bacterias & fungus (the probiotics) which help the digestive process. Naturally found prebiotics are not broken down nor absorbed in the intestines; beneficial bacterias & fungus use this fibre in fermentation. It is mainly of 3 types 1) inulin 2) oligosaccharides 3) arabinogalactans.

Inulin & oligosaccharides are short-chain polysaccharides or chain of carbs, which act at different location in the colon ensuring complete intestinal health by increasing & maintaining the population of good bacterias, fungus & yeast; & arabinogalactans are class of long, densely branched high-molecular polysaccharides.

Main sources of prebiotics: -

It is present in garlic, onions, leeks, banana, chicory root, skin of apple, beans, radish, carrots, tomato, coconut, bark of larch tree etc.

Basic clinical pharmacology of prebiotics: -

It is helpful for immune system throughout the life, helpful in fetus health, reduces risk of eczema, inhibit allergies, helpful in milk tolerance in infants, promote T helper cells which are vital for immune system, controls increased permeability of intestine, controls leaky gut, reduces inflammation of intestine, maintain health of intestines, prevent obesity, balance energy level, enhance activity of vitamins, inhibit growth of bad bacterias, help the digestive process etc.

- **Potassium: -**

It is a mineral with symbol K & atomic number 19, it is an essential mineral which body cannot prepare; it is necessary for heart, kidney & other organs to function, its low level in body is called as hypokalemia & high level is called as hyperkalemia; it is mostly present inside the cells (intracellular); normal blood range is 3.5 to 5.0 milli equivalents per/liter (mEq/L).

Main sources of potassium: -

Potassium is naturally present in figs, banana, orange, dates, raisin, broccoli, milk, chicken, sweet potato, pumpkin, spinach, watermelon, coconut water, white & black beans, potato, dried apricot, beetroot, pomegranate, almond etc.

Basic pharmacokinetics of potassium (bases on human intake in natural food products): -

It is absorbed in small intestines by passive diffusion; it is stored mostly inside the cell, little in liver, bones & red blood cells. 80 to 90% potassium is excreted in urine & 5 to 20% is excreted in stools, sweat.

Basic clinical pharmacology of potassium: -

It is a mineral belongs to electrolytes of the body; it conducts electrical impulses throughout the body & assists blood pressure, normal water balance, muscle contraction, nerves impulse, digestion, heart rhythm, maintain pH balance. It is not produced in our body so we need to consume it through eating; Kidneys maintain normal level of it in the body by excreting excessive amount of it in urine or reabsorb it if the amount is less in the body so that the body may reuse it. Its deficiency may cause weakness, low blood pressure, constipation, nausea, vomiting etc.

Its normal amount in body keeps blood pressure normal; water balance in body normal; prevents heart disease, stroke, osteoporosis, kidney stone etc.

- **Vitamin C: -**

It is also called as Ascorbic acid; it is an essential water soluble vitamin, very much needed by the body for many functions & absorption etc.

Main sources of vitamin C: -

It is present in watermelon, citrus fruit, fig, beetroot, broccoli, cauliflower, sprouts, capsicums, papaya, strawberries, spinach, green & red chillies, cabbage, leafy vegetables, tomato, cereals etc.

Basic pharmacokinetic of vitamin C (based on human intake in natural food products): -

It does not need to undergo digestion, 80 to 90% of it eaten is absorbed by intestine cell border by active transport & passive diffusion & through ion channels it enters the plasma via capillaries. It is very little stored in adrenal glands, pituitary gland, brain, eyes, ovaries, testes, liver, spleen, heart, kidneys, lungs, pancreas & muscles. All together body can store 5 grams of it & we need 200mg/day in order to maintain its normal level & uses, but old, disease person, smokers & alcoholic need more daily value. It is excreted in urine in the form of dehydroascorbic acid changed by liver & kidneys both, but unused vitamin C is excreted intact.

Basic clinical pharmacology of vitamin C: -

It prevent cough & cold, repairs tissue, acts as an enzyme for curtain neurotransmitter, important for immune function, it is a powerful antioxidant (donates electron to various enzymatic & non-enzymatic reactions); body prepares collagen with the help of vitamin c; it is also helpful in Alzheimer's, dementia, acts on iron absorption, it protects the body from oxidative damages, reduces stiffness of arteries, reduces tendency of platelets to clump each other, improves nitric oxide activity (dilatation of blood vessels) thus prevents high blood pressure & heart disease, also prevent eye disease, reduces risk of cataract, prevents the lining of lungs & prevents lung disease, it is a natural antihistamine (anti allergy), eliminates toxins from the body etc. Deficiency of it causes Scurvy disease (brown spots on skin occurs, swelling of gums, bleeding from all mucous membrane, spots are more on thighs & legs, the person looks pale, feel depressed, cannot move, loss of teeth, suppurative wounds occur.

- **Vitamin E: -**

It is fat soluble vitamin; it is a group of eight fat soluble compounds that includes four tocopherols & four tocotrienols.

Main sources of vitamin E: -

It is present in olive oil, almonds, cereals, wheat germ, sunflower oil, corn oil, soybean oil, peanuts, green leafy vegetables, cucumber etc.

Basic pharmacokinetics of vitamin E (based on human intake in natural food products): -

It is absorbed in small intestines & metabolized in liver & distributed through lymphatic system & stored in fat droplets of adipose tissue cells; it is mainly excreted in stool, little in urine & through skin.

Basic clinical pharmacology of vitamin E: -

It prevents coronary heart disease, supports immune system, prevent inflammation, promotes eye health, lowers the risk of cancer; It is a powerful anti-oxidant thus reduces UV damage of skin, nourishes & protects the skin when applied on face; also promotes hair growth.

- **Carotenoid: -**

It is a fat soluble; it is also called as tetraterpenoid; it is an organic pigment produced in plants giving them bright red, yellow, orange etc colour. It helps the plant to absorb light energy for photosynthesis; it protects our body from diseases & maintains health. It is of more than 600 types of which 50 to 60 types are eaten in food by human. It is not made by our body we depend on food source to be eaten.

Main sources of carotenoid: -

Carotenoid is present in olive oil, watermelon, tomato, kale, oranges, olive, carrot, plums, apricots, mango, sweet potato, kale, spinach, coriander, cucumber etc.

Basic pharmacokinetics of carotenoid (based on human intake in natural food products): -

It is fat-soluble; It first gets emulsified followed by solubilized in micellar then require bile salts & absorbed in intestine, little is absorbed in stomach; it is excreted in stools (research in on), it is stored in body fats and will convert the stored carotenoid into vitamin A when needed by the body and use it.

Basic clinical pharmacology of carotenoids: -

It is converted into vitamin A in our body, it is essential for vision, immune system, prevents cardiovascular disease, it helps reducing inflammation, cancers risk.

- **Vitamin A: -**

It is a fat soluble vitamin; it is group of unsaturated organic compound that includes retinol, retinal, retinoic acid & several provitamin A carotenoid. There are 2 types of vitamin A, 1) Vitamin A: - found in meat, poultry, fish & dairy products; 2) Provitamin A: - found in fruits, vegetables, plants; beta carotene is common type of provitamin A; it is an antioxidant, reduces wrinkles & repairs the skin damages; it is available in the market as tretinoin in tablets & creams to heal acne.

Main sources of vitamin A: -

It is present in watermelon, beetroot, fig, fish oil, carrot, green leafy vegetables, citrus fruit, sweet potato, spinach, kale etc.

Basic pharmacokinetic of vitamin A (based on human intake in natural food products): -

It is absorbed in jejunum mainly, little through skin; metabolism is in liver & excreted in urine & stools, it is conjugated with glucuronic acid & then changed into retinal & retinoic acid; retinoic acid is excreted in stool, mainly. It is stored primarily as palmitate in Kupffer's cells of liver, normal adult liver stores sufficient amount of it which is enough for 2 years for the body, little is stored in kidneys, lungs, adrenal glands, fats, retina; it is excreted in urine & stools.

Clinical pharmacology of vitamin A: -

it is needed by the body for vision and maintains eye health specially retina; it prevents night blindness; it helps in normal reproduction of cells thus prevents cancer; it is required for proper growth & development of embryo throughout the pregnancy period, it is good for skin, supports immune function; helps the heart, kidneys & lungs to work properly.

- **Vitamin B1 (Thiamin): -**

It is called as Thiamin also; it is a water soluble vitamin, it belongs to B-complex family, it is an essential micro nutrient which cannot be made by our body.

Main sources of vitamin B1: -

It is present in watermelon, spinach, legumes, fig, beetroot, banana, wheat germ, liver, egg, meat, dairy products, nuts, peas, fruits, vegetables, cereals, rice, breads, oats etc.

Basic pharmacokinetic of vitamin B1 (based on human intake in natural food products): -

Intestinal phosphatases hydrolyze thiamin to make it free & absorbed in duodenum, jejunum mainly through active transport in nutritional doses & passive diffusion in pharmacological doses, very little is known about its absorption; it is metabolized in liver; it is excreted in urine & stored little in liver, heart, kidney, brain, muscles.

Clinical pharmacology of vitamin B1: -

It is needed for metabolism of glucose, amino acids (proteins), lipids (fats) etc; every cell of the body require it to form ATP (adenosine triphosphate) as a fuel for energy, also it enables the body to use carbohydrates as sources of energy; also nerve cells, heart cells, muscles cell require it to function normally; its deficiency causes beri-beri heart disease, weight loss, confusion, malaise,

optic neuropathy, irritability, memory loss, delirium, muscles weakness, loss of appetite, tingling sensation in arms & legs, blurry vision, nausea, vomiting, reduce refluxes, shortness of breath etc; it is helpful to immune system; excessive intake of carbohydrates, protein, glucose (speacially in body builders, athletes etc) increases the need of vitamin B1.

- **Vitamin B2: -**

It is also called as Riboflavin, it is a water soluble vitamin, it is an essential micro nutrient, it helps many systems of the body; it is not synthesized in human body.

Main sources of vitamin B2: -

It is present in watermelon, liver, milk, fig, dairy products, nuts, egg, fish, leafy vegetables, almonds, mushroom, lean meat and beetroot.

Basic pharmacokinetic of vitamin B2 (based on human intake in natural food products): -

It is phosphorylated in the intestinal mucosa during absorption; mainly absorbed in upper gastrointestinal tract; the body absorbs little from a single dose beyond of 27mg; when excessive amount is eaten it is not absorbed; very little is known about its absorption. The conversion of it into its coenzymes takes place mainly in cells of small intestines, heart, liver, kidneys & throughout the body in many cells; it is excreted in urine & stored little in liver, heart, kidneys & in tissues of the body.

Basic clinical pharmacology of vitamin B2: -

It is needed by the body to keep skin, eyes, nerves, red blood cells healthy, it also helps adrenal gland, nerve cells, heart, brain to function; it also act in metabolism of food, amino acids (protein), fats, helps to convert carbohydrate into energy (Adenosine triphosphate formation- the energy body runs on). It plays an important role in functioning of mitochondria.

Its deficiency is called as Ariboflavinosis & causes weakness, throat swelling, soreness of mouth & tongue, cracks on skin, dermatitis, anemia, weak vision, itching & irritation in eyes, migraine.

- **Vitamin B3: -**

It is called as Niacin or Nicotinic acid; it is in 2 forms niacin & nicotinamide acid; it is water soluble vitamin; it is an essential micro nutrient; it plays a role in over 200 enzymatic reactions in the body; It is produced in the body in small amount from tryptophan which is found in protein containing food & sufficient amount of magnesium, vitamin B6 & B2 (are needed to produce it).

Main sources of vitamin B3: -

It is present in watermelon, figs, green peas, peanuts, mushroom, avocados, meat, egg, fish, milk, cereal, green vegetables, liver, chicken, coffee, potato, corn, pumpkin, tomato, almonds, spinach, enriched bread, carrots, beetroot etc.

Basic pharmacokinetic of vitamin B3 (based on human intake in natural food products): -

If eaten in natural form it is absorbed in stomach & small intestines by the process of sodium-dependent carrier-mediated diffusion in 5 to 20 minutes; if taken in therapeutic doses get absorbed by passive diffusion in small intestines. Its uptake in brain requires energy, in kidneys & red blood cells requires a carrier. It is metabolized in liver in 2 ways either is conjugated with glycine or niacin is form into nicotinamide; it is stored little in liver unbounded to enzymes. It is excreted in urine.

Basic clinical pharmacology of vitamin B3: -

It regulates lipid level in the body; it acts on carbohydrate to form energy sources for the body, it ease arthritis, boost brain function, every part of body needs it to function properly, it helps convert food into energy by aiding enzymes & cellular metabolism, it acts as an antioxidant. It

prevents heart disease. Deficiency of it causes pellagra, high blood cholesterol, memory loss, fatigue, depression, diarrhea, headache, skin problems, lesion in mouth, tiredness etc.

- **Vitamin B5 (pantothenic acid): -**

It is also called as pantothenic acid, it is water soluble vitamin, it is a micro nutrient, it is necessary for making blood cells; acts to convert eaten proteins, carbohydrate, fats into energy; it is a component of coenzyme A; it is used in synthesis of coenzyme A. (coenzyme A acts on transport of carbon atoms within the cell).

Main sources of vitamin B5: -

It is present in watermelon, beetroot, fig, meat, chicken, liver, kidney, fish, grains, milk, dairy products, legumes etc.

Basic pharmacokinetic of vitamin B5 (based on human intake in natural food products): -

It is converted into free form by intestinal enzymes & in nutritional doses it is absorbed in intestinal cells via sodium dependent active transport system in jejunum & pharmacological doses are absorbed by passive diffusion; after absorption the free form of it is now transported to erythrocytes via plasma, in cells pantothenic acid is converted into CoA, all the body tissues can convert it into CoA & ACP (acyl carrier protein), after these two complete their jobs they are degraded to form free pantothenic acid & other metabolites. It is excreted in urine & stools & little in exhaled in carbon dioxide.

Basic clinical pharmacology of vitamin B5: -

It promotes skin, hair & eyes health, proper functioning of nervous system & liver, formation of red blood cells, making of adrenal hormones, sex hormones; it is very helpful in constipation, rheumatoid arthritis, acne, allergies, asthma, baldness, colitis etc.

Its deficiency causes fatigue, nausea, vomiting, irritability, neurological weakness, numbness, abdominal cramps, sleep disturbances, hypoglycemia etc.

- **Vitamin B6: -**

It is also called as pyridoxine; it is involved in many aspects of macronutrients metabolism; it is present in many food products naturally.

Main sources of vitamin B6: -

It is present in watermelon, fig, chicken, bread, egg, vegetable, soyabean, whole grain cereals, brown rice, fish, legumes, beef, nuts, beans, liver, citrus fruits, starchy vegetables, potato, beetroot etc.

Basic pharmacokinetic of vitamin B6 (based on human intake in natural food products): -

It is absorbed in small intestine s, but before absorption a phosphate group has to be removed making vitamin B 6 in free form & absorbed by passive transport, now reaches liver via portal vein, in liver to get metabolized & flown into the blood stream it is bound with albumin & some are taken up by red blood cells, once getting in blood it can function & promote health & it is excreted mainly in urine & little is excreted in stools, it is very little stored in tissues, muscle tissues, liver, brain, kidneys, spleen.

Basic clinical pharmacology of vitamin B6: -

It is needed for proper development & function of brain in children; it is needed for neurotransmitter, histamine, haemoglobin synthesis & function. It serves as coenzyme (cofactor) for many reactions in the body, it is the master vitamin for processing amino acids & some hormones, it is needed by the body to prepare serotonin, melatonin & dopamine, it is better to intake it during treatment of tuberculosis. It supports adrenal glands to function; it acts as a

coenzyme in the breakdown & utilization of fats, carbohydrates, protein, it is important for immune system, it helps in treatment of nerve compression like carpal tunnel syndrome, premenstrual syndrome, depression, arthritis, high homocysteine level, diabetes, asthma, kidney stones etc. Its deficiency causes seborrheic dermatitis (eruption on skin), atrophic glossitis with ulceration, conjunctivitis, neuropathy, anaemia etc.

- **Folate (vitamin B9): -**

Folate is an essential micro nutrient, it is a natural form of vitamin B9, it serves many important functions of the body, it plays an important role in cell growth & formation of DNA, RNA & other genetic material & helps in treating many diseases; its name is derived from Latin word Folium, which means leaf, leafy vegetables have it in good amount; Folic acid is a synthetic form of vitamin B9.

Main sources of folate: -

It is present in watermelon, fig, dark green leafy vegetables, fruits, nuts, beans, dates, seafood, egg, dairy products, meat, chicken, legumes, beetroot, citrus fruits, broccoli, spinach, cereals etc.

Basic pharmacokinetic of folate (based on human intake in natural food products): -

Its absorption is complicated because folate present in food are of many different forms, some of which cannot be absorbed until broken down by intestinal enzymes; it is not absorbed more than 50%; dietary folate contains glutamate that need to separate it from glutamate before absorption starts; It is absorbed in duodenum & jejunum, after absorption it is converted into tetrahydrofolate (the active form of folate), then a methyl group is added to it to form methyltetrahydrofolate; now the body uses it for various functions & metabolism; the body can store folate 20-70mg in liver which is enough for 3 -6 months for the body; it gets excreted in urine & little in stools & bile.

Basic clinical pharmacology of folate: -

It is needed by the body to make DNA, RNA & other genetic material; it prevents many disease & conditions like anaemia, stroke, cardiac diseases, cancers, neurological diseases, macular degeneration (eye disease), palpitation, sores in mouth & tongue, hair fall, graying of hair. It is important in fertilization in male & female, essential during pregnancy to prevent neural tube defect in embryo (it is needed more), it protect us from free radicals & oxidation thus prevent cancers, it is essential in red blood cells formation, reduces high levels of homocysteine.

Its deficiency may cause anaemia, tiredness, palpitation, breathlessness, hairfall, neural tube defect in baby during pregnancy etc.

- **Sodium: -**

Here we are learning natural sodium, its symbol is Na & atomic no. 11; it is not produced in the body we need to take it in food sources; it is an important & essential mineral on which our body functions; it regulates blood pressure, blood volume etc.

Main sources of sodium: -

Excessive intake of sodium should be avoided; fig has very less amount of sodium in it; also vegetables & fruits have less sodium in them which is good for the body. It is present in beans, meat, fish, chicken, chilli, bread, rolls, milk, celery, beetroot, fig etc.

Basic pharmacokinetic of sodium (based on human intake in natural food products): -

It is absorbed in ileum by active sodium transport because it is impermeable & in jejunum absorption takes place via mediated active transport & depends on levels of water, bicarbonate, glucose, amino acids etc; its absorption plays an important role in the absorption of chloride, amino acids, glucose & water; similar mechanism are involved in the reabsorption of it in kidneys when its

level in the body falls. It is excreted mainly in urine, little in sweat & stools. It is stores in bones & dissolved in various body fluids.

Basic clinical pharmacology of sodium: -

It is amongst the essential electrolyte within the body, it remains in extracellular fluid (outside the cell) mainly, it carries electrical charges within the body, kidney maintain its normal level in the body, normal level is 135-145 milli-equivalent per liter (mEq/L), it is not produce in the body, it acts on muscles contraction, nerve cells, regulates blood pressure, blood volume; it takes part in every function of the body mostly, its low level in body is called as hyponatremia, it is found more in older aged, kidney disease, heart disease, hospitalized patient, this condition may cause brain edema, low blood pressure, fatigue, tiredness etc; its high level in the body is called as hypernatremia may cause increase in blood pressure, thirst, confusion, muscle twitching or spasm, seizures, weakness, nausea, loss of appetite, swelling in body etc.

- **Calcium: -**

It is natural essential mineral for the body, it is among the electrolytes of the body; its symbol is Ca & atomic no. 20.

Main sources of calcium: -

It is present in watermelon, milk, banana, cheese, fig, green leafy vegetables, soya beans, nuts, fish, meat, egg, bread, flour, yogurt, almonds, kale, soybean, spinach, beetroot etc.

Basic pharmacokinetics of calcium (based on human intake in natural food products): -

Calcium is absorbed in duodenum & upper jejunum (when calcium intake is low) by transcellular active transport process, this depends on action of calcitriol & intestinal vitamin D receptors & when calcium intake is high, absorbed by paracellular passive process throughout the length of small intestine by 3 major steps, entry across the brush border, intracellular diffusion via calcium-binding protein & extrusion; Vitamin D is necessary for absorption of calcium, also vitamin C, E, k, magnesium & exercise increases the absorption of calcium. Also the level of calcium is regulated by calcitonin released by thyroid gland it reduces calcium level in blood when it is excessive & increases the excretion of calcium via kidneys; Parathyroid hormones (PTH) released by parathyroid gland increases the blood level of calcium when body need it or calcium is less in blood & promotes reabsorption of it in kidneys (calcitonin & PTH both have opposite function). Intestines can absorb 500 to 600 mg of calcium at a time; it is mostly stored in bone tissues & teeth & excreted in stool & sweat & little in urine depended upon the level of it in blood. Also estrogen act on transport of blood calcium in bones thus women mostly suffer from osteoporosis after menopause.

Basic clinical pharmacology of calcium: -

Calcium acts on bone health, communication between brain & other parts of the body, muscles contraction, blood clotting; it is a co-factor for many enzymes, it relaxes the smooth muscles & blood vessels; it maintains heart rhythm, muscles function; it is more needed in childhood & deficiency of it in childhood may cause convulsions (seizure); Excessive level of it in blood is called as hypercalcemia & may lead to kidney stone formation, heart attack, stroke, loss of appetite, excessive urination, memory loss etc; its low level in blood is called as hypocalcemia & may lead to cramps in the body, weak bones, weak teeth, numbness, tingling etc.

Contraindication: -

Sarcoidosis, excessive level of calcium in blood, very severe constipation, kidney stones, increased activity of parathyroid gland etc. Hypersensitivity of calcium, severe cardiac diseases, hypercalcemia, hypercalciuria, severe kidney stones etc.

- **Iron: -**

It is an essential mineral for our body; its symbol is Fe & atomic no. 26; it is an important component of haemoglobin (haemoglobin binds oxygen in lungs & supply it to whole body, it is oxygen carrier).

Main sources of iron: -

It is present in watermelon, meat, dates, spinach, egg, nuts, dark leafy green vegetables, broccoli, pumpkin seeds, chicken, legumes, fish, banana, cabbage, kidney, almonds, beetroot, fig etc.

Meat is the best source of iron, it provides Fe⁺² directly which can be transported from intestine to blood stream through Fe⁺² transporter ferroportin (this binds with transferrin & delivered into tissues).

Basic pharmacokinetics of iron (based on human intake in natural food products): -

The absorption of iron is not known fully; about only 10% of iron taken in food is absorbed; it is absorbed in duodenum & upper jejunum mainly & at the end part of ileum; low pH is needed for its absorption, after absorption it get bind to transferrin (each transferrin can carry 2 atoms of iron); ceruloplasmin (protein) also helps in binding of iron; Heparin a hormone produced by liver is released when iron stores are full & inhibits iron transport & binding, thus reduces the absorption of iron; vitamin C & copper enhances iron absorption.

Storage of iron: -

Iron is stored in liver (in hepatocytes & kupffer's cells) kupffer's cells play an important role in recycling body iron, they ingest aged RBC liberate iron for it & reuse by breaking down haemoglobin. Little iron is stored in liver, heart, & kidneys in form of ferritin also little in bone marrow, spleen.

Excretion of iron: -

The body does not possess a physiological mechanism for regularly eliminating iron from the body because most of it is recycled by liver cells; iron is lost within cells, from skin & interior surface of the body (intestines, urine, breathe).

Basic clinical pharmacology of iron: -

It is an important component of Haemoglobin (haemoglobin bind oxygen in lungs & supply it to whole body); iron is beneficial for nails, hair, skin etc; it acts on blood production, its deficiency causes Anaemia (low haemoglobin level in blood) (this causes reduced in oxygen carrying capacity & supply of it); most of the iron is present in haemoglobin, it consist of one heme (iron), one protein chain (globin) this allows it to bind & load oxygen from the lungs & supply it to whole body. Unbound or free iron is highly destructive & dangerous it can trigger free radical activity which can cause cell death & destroy DNA.

- **Magnesium: -**

It is an important essential mineral; its symbol is Mg & atomic no. 12; it is a co-factor for more than 300 enzymes that regulates functions in the body. Its normal range in blood is 0.75 to 0.95 millimoles (mmol)/L.

Main sources of magnesium: -

It is present in watermelon, spinach, beetroot, meat, egg, nuts, dark leafy green vegetables, broccoli, pumpkin seeds, dates, chicken, fish, legumes, fig etc.

Basic pharmacokinetics of magnesium (based on human intake in natural food products):

It is absorbed about 20 to 50% only; it is absorbed about 40% in distal intestine when the level of it is low via passive paracellular transport & about 5% in descending colon when the level of it is high

via active transcellular transport. Vitamin D increases its absorption & also acts on its excretion in urine. It is excreted in urine & stool; it is stored in bones.

Basic clinical pharmacology of magnesium: -

It is a co-factor for more than 300 enzymes that regulates functions in the body. It act on protein synthesis, muscles & nerve function, blood glucose, control blood pressure, it is required for energy production, bone development, synthesis of DNA & RNA. It also plays a role in active transport of calcium & potassium ions, muscles contraction, normal heart rhythm etc.

- **Phosphorus: -**

It is an essential mineral; its symbol is P & atomic no. 15, it is needed for many parts & functions of the body.

Main sources of phosphorus: -

It is present in watermelon, beetroot, meat, nuts, beans, fish, chicken, dairy products, soy, grains, lentils, fig etc.

Basic pharmacokinetics of phosphorus (based on human intake in natural food products):

It is absorbed 70-85%, it is absorbed 30% in duodenum, 20% in jejunum, 35% in ileum; it is absorbed in inorganic phosphate form by 2 separate process first when the phosphorus intake is high mainly after meals by paracellular sodium independent passive diffusion pathway & second is transcellular sodium dependant carrier-mediated pathway this falls under the control of vitamin D & etc. When calcium level is too high in the body phosphorus is less absorbed, optimum calcium : phosphorus ratio is helpful in its absorption (excess of anyone decreases the absorption of both). It is stored in bones 85% & rest in tissues; it is excreted 80% in urine & rest in stools (excretion of it is a regulatory action of parathyroid hormone (PTH), vitamin D, and fibroblast).

Basic clinical pharmacology of phosphorus: -

It is present in nature combined with oxygen as phosphate. It acts on growth of teeth, bones, repairs of cells & tissues. It plays an important role in metabolism of carbohydrate, fats, protein & ATP. It works with B-complex vitamins & helps kidney function, muscles contraction, normal heart beats, nerve impulse etc.

- **Zinc: -**

It is a trace mineral; symbol is Zn & atomic no. 30; it is necessary for human body as it plays vital role in health.

Main sources of zinc: -

It is present in watermelon, meat, fish, legumes, beans, egg, dairy products, seeds, nuts, whole grains, beetroot, fig etc.

Basic pharmacokinetics of zinc (based on human intake in natural food products): -

It is absorbed 20 to 40%, its absorption depends on its concentration & is absorbed in whole intestines (jejunum has high rate of its absorption) via carrier-mediated mechanism, it is released from food as free ions during digestion. Zinc from animal sources is easily absorbed comparing to plants sources. It is present in bile & pancreatic juices which is released in duodenum & is reused by the body this is called as endogenous zinc & zinc present in food sources is called as exogenous zinc. Its absorption depends on 2 proteins- Albumin & metallophilonein. Albumin enables zinc to be transported from plasma into enterocytes. It is stored in muscles, bones mainly & little in prostate, liver, kidneys, skin, brain, lungs, heart & pancreas. It is excreted in stools 80% & rest in urine & sweat. Metallophilonein binds to zinc to make it unavailable & excrete it in stools when zinc is

excess in the body, & production of metalloprotein is reduced when zinc is less in the body to make zinc available for the body.

Basic clinical pharmacology of zinc: -

It is necessary for immune system, prevents skin diseases, heal skin diseases, helps stimulate activity of at least 100 different enzymes in the body; it is required in little amount in the body, but children, pregnant & old aged need it more. It promotes growth in children, synthesizes DNA & acts on wound healing, it is best in treating initial diarrhea & cold cough. It improves learning, memory, fertility etc. It heals acne, attention deficit hyperactivity disorder (ADHD), osteoporosis, pneumonia etc.

- **Carbohydrate: -**

It is a macronutrient needed by the body, the body receives 4 calories per 1 gram of it; carbohydrates include sugar, glycogen, starch, dextrin, fibre & cellulose that contain only oxygen, carbon & hydrogen. It is classified in simple & complex; simple carbs are sugar & complex carbs are fibre & starch which take longer to digest. It is basic source of energy for our body.

Main sources of carbohydrates: -

It is present in watermelon (little), potato, sweet potato, bread, oats, butter, white rice, whole grain rice, pasta, lentils, banana, pineapple, beetroot etc.

Basic pharmacokinetic of carbohydrate (based on human intake in natural food products):

Its digestion begins in mouth; salivary glands release saliva & salivary amylase (enzyme) which begins the process of breaking down the polysaccharides (carbohydrates) while chewing the food; now the chewed food bolus is passed in stomach through food pipe (esophagus); gastric juice like HCL, rennin etc & eaten material are churned to form chyme in the stomach; the chyme now is passed little by little down into duodenum, pancreatic amylase are released which break the polysaccharides down into disaccharide (chain of only sugars linked together); now the chyme passes to small intestine, in it enzymes called lactase, sucrase, maltase etc breakdown disaccharides into monosaccharide (single sugar) & absorbed in upper & lower intestines, through villi present in small intestine & send into liver through venous blood present in portal veins, as per bodies need it is released in the blood stream & pancreas release insulin to use it as source of energy for the body, & extra is stored is converted into glycogen by liver & stored in liver & little is stored in muscles & tissues. Liver can reconverts glycogen in to sources of energy if body lacks for other source of energy, the undigested carbohydrates reach the large intestine (colon) where it is partly broken down & digested by intestinal bacteria, the remains are excreted in stools.

Clinical pharmacology of carbohydrates: -

Carbohydrates are main sources of body energy, it helps brain, kidney, heart, muscles, central nervous system to function, it also regulates blood glucose, it acts on uses of protein as energy, breakdown of fatty acids & prevent ketosis. So it is an instant energy provider for the body & best for pre & post workout. If we eat less carbohydrate it may lead to hypoglycemia, ketosis, frequent urination, fatigue, dizziness, headache, constipation, bad breath, dehydration etc.

Excessive intake of carbohydrates may lead to vascular disease, atherosclerosis (leads to narrowing of arteries, stroke, diabetes, obesity, fatty liver, blood pressure etc).

- **Sugar (fructose): -**

Sugar present in fig is mainly fructose; (but diabetic patients should not eat much of it).

Main sources of fructose: -

It is present in watermelon, honey, banana, apple, mango, cherry, strawberry, orange, kiwi, pears, pomegranate, apricots, carrots, yogurt, bread, lemon, lime, green beans, beetroot etc.

Basic pharmacokinetics of fructose (based on human intake in natural fruit & food products): -

Fructose digestion begins in the small intestine (more in upper jejunum) via active transport or facilitated transport (not known properly). Our body cannot absorb intact polysaccharide molecules. Therefore, if fructose is present in the form of sucrose, sucrase, an enzyme, must first break up sucrose into separate glucose and fructose components. Single fructose molecules then enter the lining of the small intestine through a special channel and exit out the other side into the bloodstream, once in the bloodstream, fructose travels with all other absorbed nutrients to the liver for metabolism and processing.

Metabolism: -

Fructose metabolism occurs entirely in the liver. Through a complicated process called fructolysis, fructose undergoes several chemical and structural changes with the help of aldolase B (an enzyme in the liver).

Extra fructose needs to be changed into glycogen by liver & stored in liver, once the storage is full in liver then liver convert it into triglycerides & triglycerides are further converted by liver into very low-density lipoprotein (VLDL) & stored in fat cells & muscles. Excessive fructose is excreted in urine.

Basic clinical pharmacology of fructose: -

Fructose has low glycemic index & results in moderate release of insulin in the blood stream relative to glucose & sucrose; fructose gives the least dental caries among other types of sugars, fructose is more sweeter than other types of sugar; it does not raises blood sugar much as glucose does, it is used as sources of energy in the body, excessive intake of it may cause fatty liver, metabolic disorder, blood pressure, increase lipids, increase in uric acid level, increase in free radicals etc.

- **Choline: -**

It is water soluble vitamin & essential nutrient, it is a constituent of lecithin; it helps in many functions of the body.

Main sources of choline: -

It is present in watermelon, egg, peanut, fish, dairy products, wheat, beetroot, spinach, beans, whole grains, fig etc.

Basic pharmacokinetics of choline (based on human intake in natural food products): -

Choline is mostly present in food in free form; it is absorbed in small intestine via transporter proteins & metabolized in liver; excessive choline is not stored but converted into phospholipids; it is changed into Trimethylamine in liver & is excreted in urine.

Basic clinical pharmacology of choline: -

It helps the nerves to develop signals. Our body makes some amount of choline, but should be consumed to avoid deficiency; it helps liver function, brain development, muscles movement, cell messenger system, DNA synthesis, nervous system, gall bladder function; it can be taken in pregnancy because it prevents neural tube defect. It aids in fats & cholesterol metabolism & prevent excessive fat building in liver.

- **Copper: -**

It is an essential micronutrient mineral; its symbol is Cu & atomic no. 29; there are lot of health benefits of it; it is needed in little amount in the body.

Main sources of copper: -

It is present in beetroot leaves, watermelon, spirulina (water-plant), nuts, seeds, lobster, leafy green vegetables, guava, grapes, green olive, kiwi, mango, pineapple, pomegranate, egg etc.

Basic pharmacokinetics of copper (based on human intake in natural food products): -

It is absorbed 30 to 50%; it is absorbed easily than other minerals, its absorption depends on the copper present in the body, when the intake of it is less, absorption is increased & when intake is more absorption is less, it is mainly absorbed in small intestines & little in stomach via carrier-mediated process; its absorption is influenced by amino acids, vitamin C & other dietary factors. After absorption it is bound primarily to albumin, peptide & amino acids & transported to liver. Copper is secreted into plasma as a complex with ceruloplasmin. It is mainly stored in liver little in brain, heart & kidneys; it is excreted mainly in bile & little in urine.

Basic clinical pharmacology of copper: -

Together with iron it enables the body to form RBC; it helps to maintain health of bones, blood vessels, nerves & immune system; it also acts on iron absorption, protein metabolism, growth of body, it acts also on development of brain, heart & other organ; it is needed by the body for making ATP, collagen. Excessive of it may cause Wilson's disease.

Deficiency of copper: -

It is very rare; but may cause cardiovascular disease, genetic defects, inflammation of optic nerve etc.

- **Dietary fibre: -**

It is an eatable part of vegetables & fruit; our body cannot digest it just passes the small intestines & colon & excrete in stools; it is of two types 1) soluble fibre 2) insoluble fibre.

Soluble fibre dissolve in water & form a gel like material & helps in controlling blood cholesterol & blood glucose; it is found in apple, carrot, barley, oats, peas, beans, watermelon, fig etc.

Insoluble fibre do not dissolve & promotes excretion & increase bulk of the stool thus relief constipation & helps in elimination of toxins also. It is found in wheat flour, beans, cauliflower, potato, green beans, watermelon etc.

Watermelon & fig has both types of fibres in it, this is the reason it is helpful in constipation conditions, it can eaten in pregnancy to relief constipation and get other benefits of it also.

Basic pharmacokinetics of dietary fibre (based on human intake in natural food products):

Soluble fibres get dissolve in water & become a gelatinous substance; do not get digested; it helps to slow the digestion & help the body to absorb vital nutrient from eaten food.

Insoluble fibres do not dissolve in water but remain in fibrous form, and do not get digested; it helps the food pass through the digestive system and increase the bulk of stool & eliminate toxins also.

Basic clinical pharmacology of dietary fibre: -

It helps in slow down the digestive process thus gives a good control in blood glucose, improves insulin sensitivity, reduces risk of diabetes, maintains weight, helpful in obesity, reduces blood pressure, reduces cholesterol, reduces inflammation, reduces risk of heart disease, relieves constipation thus helpful in piles, fistula & other rectal disorders & disease, improves bowel movement thus improves bowel health, slowdowns the digestion thus improves quality of digestion, reduces risk of many types of cancer.

Its uses: -

It is used as a medicine, in milk shake, sweets, juices, custard, cooked in food items, ice creams, jams, pudding, eaten raw (fresh or dried), good for patients suffering from typhoid, dengue, malaria, cold cough, weakness, recovery stage, body building, piles, arthritis etc.

- **Natural Chloride: -**

It is mostly found in table salt, sea water, vegetables & some fruits. It is important mineral which maintaining water balance & it essential component of gastric juices, we get is from food main in the form of sodium chloride (salt). It is among electrolytes of the body. It is mostly present in body fluids. It act on electric impulses of the body, it combines with hydrogen in the stomach to form hydrochloric acid (HCL) a power digestive enzyme. It is also a by-product of the reactions between chlorine & electrolytes such potassium, sodium, magnesium which are essential for human metabolism.

Main sources of chloride: -

Tomato, celery, olives, cabbage, broccoli, radish, cauliflower, potato, pepper, eggplant etc.

- **Selenium: -**

It is an essential trace mineral, it is micro nutrient helpful to our body; its symbol is Se & atomic no. 34.

Main sources of selenium: -

It is present in watermelon, fish, nuts, beef, chicken, mushroom, egg, grains, garlic etc.

Basic pharmacokinetics of selenium (based on human intake in natural food products): -

It is mainly absorbed in duodenum & proximal jejunum by active transport process; Dietary selenium is in 2 forms organic (selenoimethionine) it is 90% absorbed & inorganic (selenite) it is 50% absorbed; after absorption it is send in liver via portal veins, liver turns it into selenite & then is bound with selenoproteins & send into blood stream, gets in RBC, muscles, tissues etc; it is not distributed evenly in the body, liver has more of it; Vitamin E & other vitamins increases its absorption & both work as an anti-oxidant. Natural selenium remains in the body for less than 24 hours; it is stored in amino acid in skeletal muscles, little in liver, kidneys & pancreas; it is primarily excreted in urine, stool & expired in air via lungs very little in sweat & semen.

Basic clinical pharmacology of selenium: -

It is important for many body functions, immune system, fertility (both male & female); it contributes in thyroid hormone metabolism, DNA synthesis; it protects the body from oxidative damages & infection, it is found in tissues, skeletal muscles; it helps testies & seminal vesicles in their function; it reduces the risk of miscarriages, liver disease, cancer, asthma, cardio vascular disease; deficiency of it causes pain in muscles & joints, weaken the hair, nails, white spots on nails are found etc.

- **Lutein & zeaxanthin: -**

Both are important carotenoid found in nature, they are related with beta carotene & vitamin A, they gives plants, fruits & vegetables yellow or red colour, they are absorbed best in human when taken with high-fat meal because it needs bile for digestion. Both are colour pigment found in human eye (macula & retina) they get deposited in macula & retina thus prevents many diseases of eyes.

Main sources of both: -

They are present in carrot, broccoli, kale, spinach, grapes, pumpkin, yellow vegetable, egg yolk, green leafy vegetable, orange, kiwi, corn etc.

Basic pharmacokinetics of both (based on human intake in natural food products): -

They are absorbed with the help of bile by mucosa of small intestine via passive diffusion & send to the liver via lymphatic system & in liver it is incorporated into low density & high density lipo proteins & transported to target tissues (retina etc) by specific lutein binding protein mediates the selective uptake of it. The absorption depends on the amount & sources of intake; it is 70 % absorbed; it is excreted in bile & urine & stored in liver & adipose tissues of the body.

Basic clinical pharmacology of both: -

They are powerful anti oxidant, anti diabetic, anti cancer. They prevent age-related macular degeneration, cataract, retinitis pigmentosa, retinopathy, macular degeneration, they work as light filter & protect the eye tissues from sunlight damages, they block blue light from reaching the underlying structure in the retina of eyes thus reduces the risk of light induce oxidative damage that could lead to age-related macular degeneration (AMD).

They also prevent free radicals thus prevents colon cancer, cervical cancer, lungs cancer, breast cancer, prostate cancer, vision loss, improves mental function, respirative infections, reduce high blood pressure, reduce soreness of muscles after exercise, reduce eye strain, controls diabetes, prevent heart diseases etc.

Total amino acids present in fig they in very little amount: -

- **Absorption & digestion of amino acid.**

When we eat high-protein foods, body breaks down protein into amino acids and peptides through digestive enzymes, such as pepsin & pancreas produces trypsin, chymotrypsin and other that aid in protein digestion.

Pepsin is the primary enzyme responsible for digesting protein; it acts on the protein molecules & breaks the bonds – called peptide bonds – that hold the protein molecules together. Next, these smaller chains of amino acids move in the stomach & then in small intestine where they're further broken down by enzymes released by the pancreas. Small intestine contains finger-like extensions called micro-villi. These structures enhance its ability to absorb dietary nutrients. Now the semi digested material pass through brush border and baso-lateral membranes of small intestine & di-tripeptides are absorbed by passive transport (facilitated or simple diffusion) or active transport (Na⁺ or H⁺ co-transporters) pathways. Di and tripeptides are more efficiently absorbed than free amino acids which in turns are better absorbed than oligopeptides. They're released into the bloodstream and used for various biochemical reactions.

Each amino acid has a different role in the human body. Upon absorption, some amino acids are incorporated into a new protein. Some fuel your muscles and support tissue repair. Others are used as a source of energy.

Tryptophan and tyrosine, for example, promote brain health. These amino acids support the production of neurotransmitters, leading to increased alertness and optimum nerve responses. Tryptophan also assists with serotonin production, lifting your mood and keeping depression at bay.

Phenylalanine serves as a precursor to melatonin, epinephrine, dopamine and other chemicals that regulate your mood and bodily functions. Methionine helps your body absorb selenium and zinc, two minerals that promote overall health. Some amino acids, such as isoleucine, play a vital role in hemoglobin production and glucose metabolism.

- **Tryptophan: -**

It is an amino acids (protein) that is useful in bio-synthesis of protein; it is essential in human because body cannot make it); it is a precursor of neuro-transmitter serotonin, melatonin, vitamin B3; it is a sedative also.

Main sources of tryptophan: -

Salmon oil, egg, spinach, milk, seeds, fenugreek seed, soy products, nuts, fish, meat, wheat, banana etc.

Basic pharmacokinetics of tryptophan (based on human intake in natural food products): -

It is absorbed in small intestine & reached the blood circulation, it passes the blood brain barrier & in brain cells it is metabolized into indolamine neuro-transmitter, niacin, a common example of indolamine is serotonin derivative from tryptophan. Tryptophan is converted into serotonin in the brain & body; it is believed that tryptophan supplements should be taken with carbidopa, which blocks the blood brain barrier. (Serotonin (5HTP) 5 hydroxytryptamine, is a monoamine neuro-transmitter. It contributes in feelings of well-being, happiness, reward, learning, memory, many physiological functions).

In the pathway of tryptophan/serotonin, melatonin hormone is produced. Melatonin regulates sleep-wake cycle. It is primarily released by pineal gland in brain. It controls circadian (daily clock) rhythms.

Pineal gland releases it at night more & very little in day light. It improves immune system function. Natural sources of melatonin are tomato, pomegranate, olive, grapes, broccoli, cucumber, barley, seeds, nuts etc.

Fructose malabsorption causes improper absorption of tryptophan in intestine thus leading to low level of it & may cause depression.

Basic clinical pharmacology of tryptophan: -

It is necessary for normal growth of infants; nitrogen balance in adults, it aids in sleep pattern, mood. It is necessary for melatonin & serotonin formation in body, it enhances mental & emotional well being, manages pain tolerance, weight etc. it also helps in build muscle tissue, essential for vitamin B3 production, relives insomnia, reduces anxiety, depression, migraine, OCD, helps immune system, reduces cardiac spasms, improves sleep patter etc.

- **Threonine: -**

It is an amino acid used in biosynthesis of proteins; it is an essential amino acid important for tooth enamel, collagen, elastin, nervous system, fats metabolism, it prevents fats buildup in liver, useful in intestinal disorders, anxiety, and depression.

Main sources of threonine: -

Cheese, chicken, fish, meat, lentil, black seed, nuts, soy etc.

Basic clinical pharmacology of threonine: -

It is useful in nervous system disorders, multiple sclerosis, spinal spasticity, makes bones, joints, tendons, ligament stronger, it helps the immune system, promotes heart health.

- **Isoleucine: -**

It is an amino acid that is used in the biosynthesis of proteins, it is an essential amino acid means the body cannot make it & we depend on food sources, it plays & helps many functions of the body.

Main sources of isoleucine: -

Meat, mutton, fish, cheese, egg, seeds, nuts, soybeans, milk, legumes, fenugreek seed etc.

Basic pharmacokinetics of isoleucine (based on human intake in natural food products): -

It is absorbed in small intestine by sodium-dependant active transport. It is metabolized in liver.

Basic clinical pharmacology of isoleucine: -

It promotes glucose consumption & uptake, it is anti-catabolic, enhances athletic performance & best for pre-workout, it acts on wound healing, detox of nitrogenous waste in the body, stimulates immune system, promotes secretion of many hormones, helps in hemoglobin formation, regulating blood glucose, energy in the body, built muscles, helpful to brain for its function.

- **Leucine: -**

It is branched chain amino acid (BCAA) it is ketogenic amino acid; it is necessary when we do exercise, it stimulates protein synthesis & assists in muscle building.

Main sources of leucine: -

Cheese, soybean, meat, nuts, chicken, seeds, fish, seafood, beans.

Basic clinical pharmacology of leucine: -

It helps regulate blood glucose, promotes growth, recovers the muscles & bone tissues, acts on production of growth hormones, repairs the tissues, essential for muscle building, it burns fats, controls obesity, promotes lean muscles growth.

- **Lysine: -**

It is an essential amino acid, which our body cannot prepare and we need to eat it from food sources. It necessary for many body functions, acts in building blocks of protein (muscles).

Main sources of lysine: -

Red meat, chicken, egg, fish, beans, lentils, wheat germ, nuts, soybeans, spirulina, fenugreek seed, shrimp, pumpkin seed, tuna, cheese, milk etc.

Basic pharmacokinetics of lysine (based on human intake in natural food products): -

It is absorbed from the lumen of the small intestine into the enterocytes by active transport, it undergo first pass metabolism in liver & is metabolized in liver.

Basic clinical pharmacology of lysine: -

It helps the body in tissue growth, repair muscles injury, promote collagen formation, help the body to produce enzymes, antibodies, hormones, supports immune system, its deficiency causes fatigue, irritability, nausea, hair loss, anorexia, inhibited growth, anemia, problems with reproductive system, it is very helpful in treating cold sores (herpes), control blood pressure, diabetes, osteoporosis, helps athletes performance, helpful in treating cancers, reduces anxiety, increase absorption of calcium, improves digestion & prevent leaky gut, helpful in pancreatitis.

- **Methionine: -**

It is a sulfur containing amino acid; it is essential; it plays a critical role in the metabolism & health; it act on normal cell functioning, growth & repair. It is also a chelating agent for heavy metals; due to its sulfur contain it is helpful in hair, nail health & growth & good for skin health; it reduces cholesterol by increase the production of lecithin in liver & reduces fats formation in liver, also protects kidneys, liver from hepatotoxins, it is an antioxidant. It is absorbed in lumen of small intestines into enterocytes by active transport & metabolized in liver.

Main sources of methionine: -

Meat, mutton, fish, chicken, cheese, egg, beans, milk, nuts, shellfish etc.

- **Cystine: -**

It is the oxidized dimer form of amino acid, it is nonessential; the body uses it to produce taurine & other amino acids; it is a sulfur containing amino acid; our body uses vitamin B6 with the help of

cystine; it heals burns, wounds, bronchitis, assist in supply of insulin, it increase level of glutathione in liver, lungs, kidneys & bone marrow. It is anti aging, anti inflammatory, anti arthritis, anti rheumatoid arthritis.

Main sources of cystine: -

Meat, egg, milk, garlic, onion, broccoli, oats, wheat germ, lentils etc.

- **Phenylalanine: -**

It is an aromatic essential amino acid in human; it plays a key role in biosynthesis of other amino acids; it is important in the structure & function of many proteins & enzymes. It is precursor of melanin, dopamine, noradrenalin hormone, thyroxin hormone. It is converted in tyrosine & used in biosynthesis of dopamine & noradrenalin. It improves memory, reduces pain of hunger; it is anti depressant; it is also a building block protein; it is useful in vitiligo, depression, ADHA, parkinson's, multiple sclerosis, pain, osteoarthritis, rheumatoid arthritis, fat burn & helpful in alcohol withdrawal symptoms.

Main sources of phenylalanine: -

Pumpkin seed, nuts, seeds, soy, meat, fish, chicken, egg, beans, milk etc.

- **Tyrosine: -**

It is a nonessential amino acid; it is also called as 4-hydroxyphenylalanine; it is useful in cell synthesis of protein; it is a building block protein; body prepares it from phenylalanine. It is a precursor & used to produce noradrenalin, dopamine, & thyroxin & melanin hormones. It reduces stress, improves memory, it promotes growth, mental health, skin health, fat burn. It acts as a mood elevator, anti depressant, improves memory, mental alertness, its deficiency can cause hypothyroidism leading to low blood pressure, low body temperature (hypothermia), stress, fatigue, narcolepsy; it helps thyroid gland, adrenal gland, pituitary gland to function properly. It is absorbed in small intestine by sodium-dependant active transport; after absorption it reaches the blood & crosses the blood brain barrier (BBB) & enters the brain cells & gets metabolized into catecholamine (noradrenalin). Human body regulates it amount by eating it by food sources & making inside the body (nonessential). The body does not store it much for later uses.

Main sources of tyrosine: -

Meat, fish, egg, milk, nuts, beans, oats, wheat, black seeds etc.

Dopamine: -

It regulates reward & pleasure centers in brain; it is a chemical important for memory, motor skills & etc.

Nor-adrenaline & adrenaline: -

These hormones are responsible for fight & flight response in stressful situation & also controls many functions of the body; it is secreted by adrenal glands.

Thyroxin: -

It is secreted by thyroid gland; it regulates metabolism, blood pressure, digestion, energy etc.

Melanin: -

It is pigmented hormone, gives our skin, hair, eye their colour; dark skinned people have more melanin in their skin than light skin people (depend on exposure to sunlight).

- **Valine: -**

It is an essential nutrient for vertebrates, biosynthesis of protein; it is an aliphatic & extremely hydrophobic essential amino acid; it is branched chain of amino acid (BCAA); it is important for growth, repair, blood glucose regulation, for energy; it stimulates CNS, proper mental function.

Main sources of valine: -

Cheese, soy, beans, nuts, fish, meat, chicken, mushroom, seeds, nuts, whole grains etc.

- **Histidine: -**

It is an amino acid used in biosynthesis of protein; it is semi essential amino acid, needed by human for production of histamine & also for growth & tissue repair, it is helpful in maintaining myelin sheaths that covers the nerves & protects the nerves.

Main sources of histidine: -

Meat, mutton, fish, milk, egg, seeds, nuts, chicken, cheese, soy, beans, whole grains, fenugreek seeds.

Basic pharmacokinetics of histidine (based on human intake in natural food products): -

It is absorbed in small intestine via active transport requiring the presence of sodium.

Basic clinical pharmacology of histidine: -

It plays many roles in immunity, gastric secretion & sexual functions. It is also required for blood cell formation & protects tissues against damage of radiation & heavy metals. It keeps normal pH of 7 in the body, useful in rheumatoid arthritis, allergy, ulcer & anemia caused by kidney failure or dialysis. It is an antioxidant, anti inflammatory, reduces cholesterol.

- **Arginine: -**

It is among conditional essential amino acid the body needs to function properly; it is made in liver; it plays an important role in building protein thus helpful in body building.

Main sources of arginine: -

Chicken, pumpkin seeds, spirulina, dairy products, red meat, fish, egg etc.

Basic pharmacokinetics of arginine (based on human intake in natural food products): -

It is absorbed in jejunum mainly from oral diet.

Basic clinical pharmacology of arginine: -

It releases nitric oxide in the blood & nitric oxide dilates the blood vessels thus increases the blood supply & controls high blood pressure, it improves erection, builds muscles etc. it also act on release of growth hormone, insulin & other substances in the body. It also improves heart health, athletes performance, stimulates immune system; citrulline present in watermelon is converted into arginine in kidneys, please refer lesson on watermelon.

- **Alanine: -**

It is a non essential amino acids that is present in blood plasma in its free state in high levels; it is involved in sugar & acid metabolism, protein synthesis, it increases immunity, provides energy for muscles tissues, brain & CNS, it act on tryptophan, vitamin B6 metabolism; it is an important sources of energy for muscles; it helps the body to convert simple sugar (glucose) into energy; it is produced in the body. It increases exercise capacity; reduces muscle fatigue, boost immunity, it is antioxidant; anti aging; increases muscle growth; ideal pre & post workout, reduce blood sugar, prevent liver disease, helps the liver to eliminate toxins, improves CNS functioning, helpful in benign prostate hypertrophy. It is digested in small intestine; it is converted into pyruvic acid by alanine aminotransferase-1; during fasting condition alanine derived from protein breakdown is

converted into pyruvate & used to synthesis glucose by gluconeogenesis in liver, it is excreted in urine via urea cycle. It is stored little in skeletal muscles.

Main sources of alanine: -

Meat, fish, egg, milk, aleovera, honey, black seeds, nuts etc.

- **Aspartic acid: -**

It is a non essential amino acid; it is over all negatively charged & plays an important role in synthesis of other amino acid, citric acid & urea cycles; it is found in animals, plants, sugarcane, sugarbeet. It may be a neurotransmitter; it strengthens the muscles, improves heart function, helps in maintaining mental health, reduces tiredness, improves athletic performance, increases muscle size, reduces depression & fatigue. It is absorbed in small intestine by active transport.

Main sources of aspartic acid: -

Meat, oysters, seeds, oats, avocado, sugar beet, milk, egg, nuts, cereals etc.

- **Glutamic acid: -**

It is a nonessential amino acid. It is an excitatory neuro-transmitter; it is necessary for biosynthesis of proteins; body uses it for several key functions within the body like making other neuro-transmitters such as GABA; it promotes brain health, muscles health, intelligence, mood & mental alertness. It is called as chemical messenger. It plays an important role in body's disposal of excessive waste like nitrogen. It is absorbed in lumen of small intestine into enterocytes by active transport & excreted in urine mainly. It is almost about 2 kilo grams, storage in natural form in brain, kidneys, liver, muscles etc.

Main sources of glutamic acid: -

Meat, chicken, fish, egg, milk, wheat, mushroom, soy, broccoli, walnut, peas etc.

- **Glycine: -**

It is a nonessential amino acid that body needs for growth & maintainance of tissue & need to prepare hormones & enzymes. It is inhibitory neurotransmitter. It helps in preparing glutathione (a powerful antioxidant & reduces free radicals, delay aging). It is helpful in preparing of creatine (provides energy to muscles to perform exercise etc & acts on muscle contraction), beneficial for brain health, bone health, alzheimer's, schizophrenia, sleep disorder, stroke, burns, protects kidney & liver from harmful side effects of drugs used after organ transplant, heals wound & ulcers, it is anti inflammatory, improves skin health.

Main sources of glycine: -

Meat, fish, milk, legumes etc.

- **Proline: -**

It is a protein-genic amino acid used in biosynthesis of proteins. It heals cartilages, cushion joints, tendons, ligament, heart muscles, connective tissues & helps in formation of collagen.

Main sources of proline: -

Soy, pumpkin seed, lentils, black beans, quinoa etc.

- **Serine:-**

It is a nonessential amino acid, important for synthesis of protein, fats metabolism, muscle growth, immune system; it is a precursor of many amino acids, helpful in enzyme catalyze its reaction, overall health, physical & mental health.

Main sources of serine: -

Soybean, egg, lentils, meat, fish, nuts, almonds, walnut etc.

FIG			NUTRITION FACTS			Dr. A FOOD & ME		
PRINCIPLE			NUTRIENT VALUE			PERCENT OF RDA		
CALORIES	47 Kcal	2%						
CARBOHYDRATES	12.3 g	4%						
PROTEIN	0.5 g	1%						
TOTAL FAT	0.2 g	0%						
CHOLESTEROL	0 mg	0%						
DIETARY FIBER	1.9 g	7%						
SUGARS	10.4 g							
SODIUM	1MG	0%						
POTASSIUM	149 MG	4%						
			MINERALS					
			CALCIUM	22.4MG	2%			
			IRON	0.2MG	1%			
			MAGNESIUM	10.9MG	3%			
			PHOSPHORUS	9.0MG	1%			
			POTASSIUM	149MG	4%			
			SODIUM	0.6MG	0%			
			ZINC	0.1MG	1%			
			COPPER	0.0MG	2%			
			MANGANESE	0.1MG	4%			
			SELENIUM	0.1MCG	0%			
VITAMINS								
VITAMIN A	90.9IU	2%	FOLATE	3.8MCG	1%			
VITAMIN C	1.3MG	2%	VITAMIN B	0.0MCG	0%			
VITAMIN E (ALPHA TOCOPHEROL)	0.1MG	0%	PANTOTHENIC ACID	0.2MG	2%			
VITAMIN K	3.0MCG	4%	CHOLINE	3.0MG				
VITAMIN B6	0.1MG	4%						

● **Reference/research: -**

- 1) www.britannica.com/plant/olive-plant
- 2) Origin and History of the Olive | IntechOpen
- 3) www.intechopen.com/books/origin-and-history-of-the-olive
- 4) <https://www.healthline.com/nutrition/11-proven-benefits-of-olive-oil>
- 5) www.tib-e-nabi-for-you.com
- 6) Hort, Sir Arthur (1916). Theophrastus Enquiry into Plants. William Heinemann. p. 107.
- 7) Abdelhafiz and Muhamad, 2008
- 8) A.T. Abdelhafiz, J.A. Muhamad Midcycle pericoital intravaginal bee honey and royal jelly for male factor infertility
- 9) Int. J. Gynaecol. Obstet., 101 (2) (2008), pp. 146-149
- 10) ArticleDownload PDF CrossRefView Record in ScopusGoogle Scholar
- 11) Ahmad, 2016
- 12) K. Ahmad Update on pediatric cough
- 13) Lung, 194 (2016), pp. 9-14
- 14) CrossRefView Record in ScopusGoogle Scholar
- 15) Akan and Garip, 2011
- 16) Z. Akan, A. Garip Protective role of quercetin: antioxidants may protect cancer cells from apoptosis and enhance cell durability
- 17) WebmedCentral, 2 (1) (2011)
- 18) WMC001504
- 19) Google Scholar
- 20) "beet". def. 1 and 2. also "beet-root". Oxford English Dictionary Second Edition on CD-ROM (v. 4.0) © Oxford University Press 2009
- 21) 1771 illustration from Trew, C.J., Plantae selectae quarum imagines ad exemplaria naturalia Londini, in hortis curiosorum nutrit, vol. 8: t. 73 (1771), drawing by G.D. Ehret
- 22) ^ "The Plant List".
- 23) ^ Jump up to:^{a b c d} The Fig: its History, Culture, and Curing, Gustavus A. Eisen, Washington, Govt. print. off., 1901
- 24) ^ RHS A-Z encyclopedia of garden plants. United Kingdom: Dorling Kindersley. 2008. p. 1136. ISBN 978-1405332965.
- 25) ^ Flora of China Ficus carica Linnaeus, 1753. 无花果 wu hua guo
- 26) ^ "Biota of North America Program 2014 county distribution map".
- 27) ^ T.F. Hoad, The Concise Oxford Dictionary of English Etymology, Oxford University Press, 1986, page 171a.
- 28) ^ Condit, Ira J. (1947) The Fig; Chronica Botanica Co., Waltham, Massachusetts, USA.
- 29) ^ Jump up to:^{a b} Wayne's Word: Sex Determination & Life Cycle in Ficus carica Archived 2009-09-02 at the Wayback Machine
- 30) ^ Jump up to:^{a b} "Fig, Ficus carica". Purdue University: Horticulture & Landscape Architecture. Retrieved December 6, 2014.
- 31) ^ Shannon Wolfe, "Carnivorous Figs: The Relationship Between Wasps and Figs", University of California Master Gardener Program of Contra Costa County, 27 August 2014.

- 32) ^ Louise Ferguson and Carlos H. Crisosto, "The Fig: Overview of an Ancient Fruit", in HortScience, August 2007, Vol. 42, No. 5, pages 1083-7.
- 33) ^ Jump up to:^a ^b Blackburne-Maze, Peter (2003). Fruit: an Illustrated History. Buffalo, New York: Firefly Books. pp. 209–11. ISBN 9781552977804. Retrieved March 22, 2017.
- 34) ^ Kislev et al. (2006a, b), Lev-Yadun et al. (2006)
- 35) ^ Jump up to:^a ^b Leroi, Armand Marie (2014). The Lagoon: How Aristotle Invented Science. Bloomsbury. pp. 244–247. ISBN 978-1-4088-3622-4.
- 36) ^ Mary Beard (2013). Confronting the Classics. Liveright Publishing Corporation, a division of W.W. Norton & Company. p. 128.
- 37) ^ Cassius Dio. Roman History 56.30.
- 38) ^ Mary Beard (2013). Confronting the Classics. Liveright Publishing Corporation, a division of W.W. Norton & Company. p. 131.
- 39) ^ Jump up to:^a ^b Roeding, George C. (1903) The Smyrna Fig: At Home and Abroad; published by the author, Fresno, CA, USA
- 40) ^ Jump up to:^a ^b Storey, W.B, Enderud, J.E., Saleeb, W.F., & Mauer, E.M. (1977) The Fig, *Ficus carica* Linnaeus: Its Biology, History, Culture, and Utilization, Vol. 13 #2,3,4; Jurupa Mountains Cultural Center, Riverside, CA, USA
- 41) ^ Anne Rowlands (2012-10-01). "Growing Fig Trees in Cooler Climates". conngardener.com. Retrieved 2018-04-09.
- 42) ^ Jump up to:^a ^b Fernanda Santos (2010-02-15). "Winter Coats No Longer the Fashion for Fig Trees". The New York Times. Retrieved 2018-04-09.
- 43) ^ Hal Klein (2014-12-25). "Why Bury Fig Trees? A Curious Tradition Preserves a Taste of Italy". Npr.org. Retrieved 2018-04-09.
- 44) ^ "RHS Plant Selector – *Ficus carica* 'Brown Turkey'". Retrieved 20 June 2013.
- 45) ^ "RHS Plantfinder - *Ficus carica* 'Ice Crystal'". Retrieved 27 February 2018.
- 46) ^ "AGM Plants - Ornamental" (PDF). Royal Horticultural Society. July 2017. p. 39. Retrieved 27 February 2018.
- 47) ^ Die Feigenernte in Kraichgau und Pfalz läuft auf Hochtouren(English: The fig harvest in Kraichgau and the Palatinate is in full swing) at Rhein-Neckar-Zeitung.
- 48) ^ Christoph Seiler: Feigen aus dem eigenen Garten (English: Figs from your own garden), Stuttgart 2016, page 64.
- 49) ^ Christoph Seiler: Feigen aus dem eigenen Garten. Stuttgart 2016, pages 75 and 78.
- 50) ^ "Figs". Royal Horticultural Society. Retrieved 11 December 2016.
- 51) ^ Jump up to:^a ^b "Fig". CRFG Publications. California Rare Fruit Growers. 1996. Retrieved 2016-02-01.
- 52) ^ North American Fruit Explorers: Figs. Archived 2009-04-10 at the Wayback Machine
- 53) ^ Jump up to:^a ^b Janick, Jules & Moore, James (editors) (1975) Advances in Fruit Breeding; pgs 568-588: Figs, by Storey, W.B.; Purdue University Press, West Lafayette, IN, USA
- 54) ^ Jump up to:^a ^b Van Deynze, Allen (editor) (2008) 100 years of breeding: UC Davis Plant Breeding Program. Published by the Dean's Office, Department of Plant Sciences, & Seed Biotechnology Center, Davis, CA, USA
- 55) ^ "New fig cultivar comparison report released by UC Kearney REC". westernfarmpress.com. Archived from the original on 2009-09-06. Retrieved 2016-02-01.
- 56) ^ Jump up to:^a ^b "Raw fig production in 2017; Crops/World Regions/Production Quantity from picklists". UN Food and Agriculture Organization Corporate Statistical Database, FAOSTAT. 2018. Retrieved 2 May 2019.
- 57) ^ "Fig". BBC Good Food. Retrieved 2016-02-02.
- 58) ^ "Nutrition facts for dried figs, uncooked per 100 g". Conde Nast for the USDA National Nutrient Database, version SR-21. 2014. Retrieved 5 June 2014.
- 59) ^ Vinson (1999)
- 60) ^ Veberic R, Colaric M, Stampar F (2008). "Phenolic acids and flavonoids of fig fruit (*Ficus carica* L.) in the northern Mediterranean region". Food Chemistry. 106 (1): 153–157. doi:10.1016/j.foodchem.2007.05.061.
- 61) ^ Solomon A, Golubowicz S, Yablowicz Z, Grossman S, Bergman M, Gottlieb HE, Altman A, Kerem Z and Flaishman MA (2006). "Antioxidant activities and anthocyanin content of fresh fruits of common fig (*Ficus carica* L.)". J Agric Food Chem. 54 (20): 7717–7723. doi:10.1021/jf060497h. PMID 17002444.
- 62) ^ Landranco, Guido (2001). *Medicina popolare ta' l-imghoddi fil-gzejjer Maltin* [Popular medicine of the past in the Maltese islands] (in Maltese). Valletta, Malta: Klabb Kotba Maltin. ISBN 99909-75-97-3.
- 63) ^ Polat, Muhterem; Öztaş, Pınar; Dikilitaş, Meltem Cik; Allı, Nuran (December 2008). "Phytophotodermatitis due to *Ficus carica*". Dermatol Online J. 14 (12): 9. PMID 19265622.
- 64) ^ Son, J. H.; Jin, H.; You, H. S.; Shim, W. H.; Kim, J. M.; Kim, G. W.; Kim, H. S.; Ko, H. C.; Kim, M. B.; Kim, B. S. (February 2017). "Five Cases of Phytophotodermatitis Caused by Fig Leaves and Relevant Literature Review". Annals of Dermatology. 29 (1): 86–90. doi:10.5021/ad.2017.29.1.86. PMC 5318534. PMID 28223753.
- 65) ^ "FDA Poisonous Plant Database". U.S. Food & Drug Administration. Retrieved 11 December 2018.
- 66) ^ McGovern, Thomas W.; Barkley, Theodore M. (2000). "Botanical Dermatology". The Electronic Textbook of Dermatology. Internet Dermatology Society. Section Phytophotodermatitis. Retrieved November 29, 2018.
- 67) ^ Jump up to:^a ^b Zaynoun, S. T.; Aftimos, B. G.; Abi Ali, L.; Tenekjian, K. K.; Khalidi, U.; Kurban, A. K. (July 1984). "Ficus carica; isolation and quantification of the photoactive components". Contact Dermatitis. 11 (1): 21–25. doi:10.1111/j.1600-0536.1984.tb00164.x. PMID 6744838. Cited in McGovern and Barkley 2000, section Phytophotodermatitis.
- 68) ^ Jump up to:^a ^b Li, Jun; Tian, Yu-zeng; Sun, Bao-ya; Yang, Dan; Chen, Ji-ping; Men, Qi-ming (2011). "Analysis on Volatile Constituents in Leaves and Fruits of *Ficus carica* by GC-MS" (PDF). Chinese Herbal Medicines. 4 (1): 63–69. doi:10.3969/j.issn.1674-6384.2012.01.010.
- 69) ^ Walker, Barbara (1988). The Woman's Dictionary of Symbols and Sacred Objects. Harper One. p. 484.
- 70) ^ Eric Brandon Roberts (2012). The Parables of Jesus Christ:: A Brief Analysis. Booktango. ISBN 9781468908800.
- 71) ^ Theodor Herzl (1987). Old New Land. Translated by Lotta Levensohn. M. Wiener. ISBN 9-781-55876-160-5.
- 72) ^ "George Washington and his Letter to the Jews of Newport". Touro Synagogue.
- 73) ^ "Foods of the prophet". IslamOnline.
- 74) ^ "BBC iPlayer – In our time: the destruction of Carthage". Retrieved 20 June 2013.

- 75) ^ "sycophant (n.)". Online Etymology Dictionary.
- 76) ^ बालुरघाट में दिखा गूलर का विस्मयकारी फूल, Sep 20, 11:39 pm (Hindi version), (Translated version)
- 77) ^ Gular ka phool by Rajiv Kumar Trigarti
- 78) ^ A review of the early Miocene Mastixioid flora of the Kristina Mine at Hrádek nad Nisou in North Bohemia, The Czech Republic, January 2012 by F. Holý, Z. Kvaček and Vasilis Teodoridis - ACTA MUSEI NATIONALIS PRAGAE Series B – Historia Naturalis • vol. 68 • 2012 • no. 3–4 • pp. 53–118

● **Research: -**

SCIENCE & HADEES REGARDING FIGS: -

Nabi ﷺ said regarding figs that, "If I had to mention a fruit that descended from paradise (Jannat) I would say this is it because the paradisiacal fruits do not have pit, eat from these fruits for they prevent hemorrhoids piles and help gout" (Arthritis).

Figs are a top source of fiber, as well as potassium and vitamin B6. Fiber results in bulkier stools, which lessen the incidence of constipation, hemorrhoids and colon (large intestine) cancer. Fiber also lowers cholesterol and the risk of heart disease.

Just one fig provides 20% of the (USDA) fiber. A Harvard University study of 43,757 men 40 to 75 were found that those who got the most fiber had about half the risk of getting a heart attack as those who got the least. Furthermore, men who add just 10 grams of fiber a day to their diet (6 figs) lowered their incidence of heart disease by 30%.

Conclusion of research: -

1. Figs are from Jannah & are dry fruits of Jannah. Recite Chapter 95 (Sura) Teen In Isha Salah. It cures Piles & Arthritis. Many interesting biological activities of *F. carica* have been carried out, which can be further explored to make use of them as a healing method for the future. For example, the leaves have shown irritant activity; consequently they can be investigated against parasitic infection and ovicidal activity. The majority of the pharmacological studies which have been carried out on *F. carica* were conducted with uncharacterized crude extracts; it is difficult to produce the grades of these studies and identify the bioactive metabolites.

Phytochemical research carried out on *F. carica* has led to the isolation of few classes of plant metabolites. Most of the phytochemical works have been employed on leaves and fruits of *F. carica*, while there is little information on stem and root phenolic profiles. However the vast traditional uses and established pharmacological activities of *F. carica* point out that an enormous scope still exists for its phytochemical exploration using bioassay-guided isolation. The result of future research in the above mentioned areas will afford a persuasive support for the future clinical uses of *F. carica* in contemporary remedy.